

UPPER BLEPHAROPLASTY w/out Lid Lift Goggles POST-OP INSTRUCTIONS

- Apply cool compresses to the eyes on arrival home from surgery as follows: Place 4x4 gauze pads soaked in cool <u>Sterile Saline</u> over the eyes for first 48 hours. Place frozen peas or cool compress on top of this, 20 minutes on, 20 minutes off (while awake).
- Apply Vaseline ointment to eyelid suture lines with a Q-tip 3 times a day starting morning after surgery.
- Begin Natural Tears® eye drops 3 times a day starting the morning after surgery.
- Sleep elevated on 3-4 pillows or recliner chair for the first week to decrease swelling.
- No driving for 3 days/ No lifting or extreme exercise for 2 weeks/ No contacts in for 2 weeks.

EXPECTATIONS

- Rust colored or bloody tears are normal for the first week.
- Temporary bruising and swelling will occur during the first week.
- You may experience blurred vision and tearing for approximately 2 weeks.
- Blinking may be difficult or sluggish for the first week.
- Avoid sun or windburn for six months.
- After surgery, Dr. Gross recommends using one of our Medical Grade skin care eye creams for best maintenance care.
- Women may begin to apply make-up around the tenth day after surgery (check w/ Dr. Gross for his approval).

MEDICATIONS

- □ Doxycycline 100mg 1 tablet two times a day for 3 days / antibiotic.
- □ Lortab 7.5mg (Hydrocodone) 1 tablet every 6 hours as needed / pain.
- □ Celebrex 200mg take once a day as directed / inflammation.
- □ Ativan 1mg (Lorazepam) 1 tablet every 8 hours as needed / anxiety and sleep.
- □ Natural Tears® eye drops three times a day to lubricate eyes.
- ☐ Arnica take as directed (begin day before surgery) / swelling and bruising.
- □ Phenergan (promethazine) 1 tablet every 6 hours as needed / nausea

Follow up Appoir Primera call or Te		ions or concerns. If you have an emergency, call	l
911.	-		
I have received a	copy of the above instructions and	l agree to follow them	
X	Date:	<i>Time</i> :	
		Revii.Zi	